




# **Chapter Seven**

# **The Electronic Structure of Atoms**

<http://clipart-library.com>

# Wave Theory

## Wave

- Repeating disturbance spreading out from a defined origin
- Characterized by wavelength, frequency and amplitude

## Wavelength ( $\lambda$ )


- Distance between identical pts
- Units some form of meters

## Frequency ( $\nu$ )

- Number of waves that pass through a point in 1 second
- Units of cycles/sec or Hz

## Amplitude



- Height of wave from center point
- Intensity of wave



# Electromagnetic Radiation

## Electromagnetic Radiation

- Emission/transmission of energy
- In form of waves
- Has electrical & magnetic components
- Travels at the speed of light ( $c = 3.00 \times 10^8 \text{ m/s}$ )



Frequency ( $\nu$ , Hz) & wavelength ( $\lambda$ , nm) are linked:

$$c = \lambda \nu$$

$$\text{Hz} = \text{s}^{-1}$$

**Using the relationship  $c = \lambda\nu$ :**

What is the wavelength of an FM-radiowave with a 94.9 MHz frequency?

**A: 3.16 m**

# Max Planck's Quantum Theory

Studied energy emitted by objects

- Amount of energy emitted was directly related to wavelength at which energy was emitted

Theory: Energy must be in discrete amounts.

- Amounts were defined by  $\lambda$  (&  $\nu$  – they are related!)

$$E = h\nu = hc/\lambda$$

- Can have multiples of these discrete amounts

$$E = h\nu, E = 2h\nu, E = 3h\nu \dots$$

- **$h$  = Planck's constant =  $6.626 \times 10^{-34} \text{ J s}$**

Called the smallest amount of energy a **Quantum**.

Didn't know why, but math worked over entire spectrum

# Einstein and the Photoelectric Effect


Experiment to prove why  $E = h\nu$

- Full spectrum of light hits metal surface
- Energy transferred to electrons in metal
- Electrons break free and escape to anode
- Flow of electrons recorded with voltmeter
- **Light energy must be at or above a certain frequency to dislodge electrons**

Conclusions:

- **Light energy has wave properties:  $E = h\nu$**   
&
- **Light energy has particle properties**

Particles of light were later called  
**“photons”**



Using  $E = h\nu$  ( $h = 6.626 \times 10^{-34} \text{ Js}$ )

What is the energy of a radiowave with a frequency of 94.9 MHz? A:  $6.29 \times 10^{-26} \text{ J}$

What wavelength has an energy of  $1.00 \times 10^{-20} \text{ J}$ ?

A:  $1.99 \times 10^{-5} \text{ m}$   
Or  $19.9 \mu\text{m}$

Using  $E = h\nu$  ( $h = 6.626 \times 10^{-34} \text{ Js}$ )

What is the energy per photon and per mole of photons of violet light, with a wavelength of 415 nm?



A:  $4.79 \times 10^{-19} \text{ J/photon}$   
A:  $2.88 \times 10^5 \text{ J/mol}$

# Elemental Line Spectra

**Emission Spectra:** Pattern of radiation that is emitted when photons are removed from a substance.

## Procedure

- Add energy to an element
- Photons are emitted as a beam of light
- Separate wavelengths through a prism
- Record pattern on a photographic plate



# Continuous vs. Line Spectra


Continuous spectrum:

- Occurs when all visible light is present: white light




Line Spectrum

- Occurs when light is produced through an element
- Pattern of lines is characteristic of the element
- Can be used for identification of elements



# Elemental Line Spectra


## Bright-line Spectra



# Bohr's Hydrogen Atom

Niels Bohr (1913): Electron energy ( $E_n$ ) was quantized

- Only certain specified values allowed
- Stable levels called energy levels
- Photon absorbed/released when electron moves from 1 level to another




The energy of each stable orbit:  $E_n = -R_H/n^2$

- $n$  is the quantum number of the level
- $n$  is always an integer, 1,2,3,...etc.

Proportionality constant  $R_H$

- Rydberg constant
- **$R_H = 2.18 \times 10^{-18} \text{ J}$**

Leads to orbit description of atoms




# Energy Level Calculations

All calculations done by comparing energy levels

- Electron moves between levels
- $E = -R_H (1/n_f^2 - 1/n_i^2)$

Energy emitted or absorbed

- High to low level:
  - energy released (-)
- Low to high level:
  - energy absorbed (+)



**Ground state:** An  $e^-$ 's lowest possible energy level

**Excited state:** All other levels

Calculate the wavelength of the electron shift from<sup>14</sup>  
 $n = 4$  to  $n = 2$ . Is light emitted or absorbed?



$$E = -R_H \left( \frac{1}{n_f^2} - \frac{1}{n_i^2} \right)$$

$$R_H = 2.18 \times 10^{-18} \text{ J}$$

A:  $\lambda = 486 \text{ nm}$  Visible blue green light is emitted (neg E value)

# Modern View of the Atom: Quantum Mechanics – a very brief intro

- (Nucleus in center, protons & neutrons in nucleus)
- Electrons outside nucleus
  - located in “cloud” surrounding the nucleus
  - likely location based on probability functions
  - quantum numbers used to describe probable location



$$i\hbar \frac{\partial \Psi}{\partial t} = -\frac{\hbar^2}{2m} \nabla^2 \Psi + [V_1(x) + iV_2(x)] \Psi$$

# Quantum Numbers and Atomic Orbitals

## Atomic orbital



- A region in space with a high probability of finding an electron.
- Identified by 4 quantum numbers.

## 4 Quantum Numbers (think of it as a dorm address)


- |                                            |          |
|--------------------------------------------|----------|
| 1. Principal quantum number ( $n$ ):       | Building |
| 2. Angular momentum quantum number ( $l$ ) | Floor    |
| 3. Magnetic quantum number ( $m_l$ )       | Room #   |
| 4. Electron spin quantum number ( $m_s$ )  | Bed      |

# The Principal Quantum Number (n)

- Restricted to the positive integers: 1, 2, 3, 4, 5, 6, 7
- The shell or energy level of the orbital



- Indicates the size of the orbital
  - max distance  $e^-$  can travel from nucleus
- Integers correspond to row numbers in Periodic Table
  - row an element is in tells you the highest energy level in the ground state



# The Angular Momentum Quantum Number ( $\ell$ ) <sup>18</sup>

- Indicates orbital shape

- Designation: s, p, d or f

| level | 0 | 1 | 2 | 3 |
|-------|---|---|---|---|
| Name  | s | p | d | f |

- Designates the **subshell**


- Values range from 0 to  $n-1$
- 0-6 theoretically, but realistically 0-3
- Give rise to “Blocks” in periodic table

| Energy Level (n) | Math      | Allowed $\ell$ values | Orbitals  |
|------------------|-----------|-----------------------|-----------|
| 1                | $1-1 = 0$ | 0                     | s only    |
| 2                | $2-1 = 1$ | 0, 1                  | s & p     |
| 3                | $3-1 = 2$ | 0, 1, 2               | s, p, & d |

# Orbital Shapes = $\ell$ quantum number


## $\ell = 0$ : s orbitals

- Spherical
- **One** per energy level




## $\ell = 1$ : p orbitals

- 2 teardrops joined at center
- **Three** per energy level




## $\ell = 2$ : d orbitals

- Most are like two p orbitals along different axes
- **Five** per energy level



## $\ell = 3$ : f orbitals.

- Complicated shapes
- **Seven** per energy level



# The Magnetic Quantum Number ( $m_\ell$ ):

Determines the orientation in space of the orbitals

- “orientation” refers to proximity to axes (x, y, z)
- Integers from  $-\ell$  to  $+\ell$

Determines the number of orbitals in a subshell

- The number of possible values for  $m_\ell = 2\ell + 1$

| Orbital | $\ell$ value | Allowed $m_\ell$ values | Number of Orbitals per Energy Level |
|---------|--------------|-------------------------|-------------------------------------|
| s       | 0            | 0                       | 1                                   |
| p       | 1            | -1, 0, 1                | 3                                   |
| d       | 2            | -2, -1, 0, 1, 2         | 5                                   |
| f       | 3            | -3, -2, -1, 0, 1, 2, 3  | 7                                   |

Orbitals with same  $n$  &  $\ell$  values are “degenerate”

degenerate = same energy

(Note: In some cases there are slight energy differences)


Possible quantum numbers for an electron in a 3p orbital:

$$n = 3$$

$\ell$  can be 0 to  $3-1$  (0, 1, 2) BUT if it is a p orbital  $\ell = 1$


$m_\ell$  can be  $+\ell$  to  $-\ell = -1, 0, +1$

Since the 3p orbitals are degenerate, any of the three  $m_\ell$  values could be correct



# Electron Spin Quantum Number ( $m_s$ )

- A magnetic field is induced by the moving electric charge of an electron as it spins
  - Opposite spins cancel one another
  - No net magnetic field for the pair
  - Allows 2 electrons to occupy 1 orbital
  - Unpaired  $e^-$  lead to magnetism
- Two possible values:  **$+1/2$  and  $-1/2$**



# Quantum Numbers Summary

TABLE 7.2

Relation Between Quantum Numbers and Atomic Orbitals

| $n$ | $\ell$ | $m_\ell$          | Number<br>of Orbitals | Atomic<br>Orbital Designations                           |
|-----|--------|-------------------|-----------------------|----------------------------------------------------------|
| 1   | 0      | 0                 | 1                     | $1s$                                                     |
| 2   | 0      | 0                 | 1                     | $2s$                                                     |
|     | 1      | $-1, 0, 1$        | 3                     | $2p_x, 2p_y, 2p_z$                                       |
| 3   | 0      | 0                 | 1                     | $3s$                                                     |
|     | 1      | $-1, 0, 1$        | 3                     | $3p_x, 3p_y, 3p_z$                                       |
|     | 2      | $-2, -1, 0, 1, 2$ | 5                     | $3d_{xy}, 3d_{yz}, 3d_{xz},$<br>$3d_{x^2-y^2}, 3d_{z^2}$ |
| ⋮   | ⋮      | ⋮                 | ⋮                     | ⋮                                                        |
| ⋮   | ⋮      | ⋮                 | ⋮                     | ⋮                                                        |
| ⋮   | ⋮      | ⋮                 | ⋮                     | ⋮                                                        |

# Quantum Numbers & the Periodic Table

- **Principle quantum number,  $n$** 
  - Row number of periodic table, values of 1-7
- **Angular momentum quantum number,  $\ell$** 
  - Specific area of periodic table, spdf “blocks”
- **Can follow the periodic table to fill  $e^-$  configuration**
- **Can use location on Periodic Table to determine where  $e^-$  configuration will end**

**Electrons in the outermost energy level are the valence electrons.**

|    |    |    |    |
|----|----|----|----|
| 1s |    |    | 1s |
| 2s |    |    | 2p |
| 3s |    |    | 3p |
| 4s | 3d |    | 4p |
| 5s | 4d |    | 5p |
| 6s | 5d |    | 6p |
| 7s | 6d |    | 7p |
|    |    | 4f |    |
|    |    | 5f |    |

A possible set of quantum numbers for the last electron added to complete an atom of selenium would be:

$n$ :

$l$ :

$m_l$ :

$m_s$ :

# Electron Configuration:

## Finding a home for each electron


The energy of an electron is defined by both  $n$  &  $\ell$

- Principle shells (size)

- $n = 1, 2, 3, 4$  or  $5$

- Subshells (shape)


- $\ell = 0, 1, 2$ , or  $3$
- $n$  determines number of subshells
- $s, p, d, f$  orbitals
- Shielding impacts relative energies



# Rules & Principles Governing e<sup>-</sup> Configurations


## Pauli Exclusion Principle:

- No 2 e<sup>-</sup> in an atom can have the same set of 4 quantum #s
  - If in the same orbital, e<sup>-</sup> must have opposite spins



## Hund's rule:


- Electrons in the same subshell occupy degenerate orbitals singly, before pairing
  - Degenerate = same energy



Ex: Oxygen, O    Z = 8


## The Aufbau Principle:

- In general, each successive electron added to an atom occupies the lowest energy orbital available
  - There are some exceptions




# Orbital Filling in Multi-electron Atoms

- Fill low to high energy
  - 2 electrons per orbital
  - Shielding impacts the energy of orbitals
  - Use chart to account for overlap of n values
- $1s < 2s < 2p < 3s < 3p < 4s < 3d < 4p < 5s < 4d < 5p < 6s$
- Format: spdf or orbital notation
  - Ends when a home is found for each electron



# Electron Configuration con't

- Defines the orbital (“home”) for each electron
- # electrons = atomic number (Z) of atom (if neutral)
  - Max 2 electrons per orbital



## Orbital Diagrams



- Energy increases from bottom to top
  - Higher energy levels at top
- Boxes or lines represent orbitals
  - # lines at one level = # degenerate orbitals
- Arrows ( $\uparrow\downarrow$ ) represent  $e^-$
- 2  $e^-$  allowed per orbital
  - one arrow up & one down to show the different spins

# Formats for Electron Configurations

## spdf Notation

- Front number = energy level
- Letter = type of orbital (s, p, d, or f)
  - Degenerate orbitals are combined together
- Superscript = # electrons in that type of orbital
  - Degenerate orbitals are combined, so the superscript can be more than 2 if it is a p, d, or f orbital (p max 6, d max 10, f max 14)


Examples:



## Orbital Notation

- Number = energy level
- Letter = type of orbital
  - Degenerate orbitals are NOT combined
- Arrows = electrons
  - Put one  $e^-$  in each degenerate orbital before pairing
  - If 2  $e^-$  in one orbital, one arrow must be up, the other down

Example:



# Writing Electron Configurations

Sulfur:

Vanadium:

# Writing Electron Configurations for Ions

**Remove Electrons from Highest Energy Level First**

Magnesium ion ( $\text{Mg}^{2+}$ ):

Fluorine ion ( $\text{F}^-$ ):

Manganese (II) ion ( $\text{Mn}^{2+}$ ):

# Noble Gas Configuration

- Abbreviation of Electron Configuration (ex: Na: [Ne]3s<sup>1</sup>)
- Noble gas symbol replaces the portion of the e<sup>-</sup> config. that is identical to the e<sup>-</sup> config. of the noble gas.
- Always use the largest noble gas that is smaller than the element
- Can use for either spdf or orbital notation
- Will always start with an s orbital – just not 1s.
- ex: Arsenic  
regular configuration: 1s<sup>2</sup>2s<sup>2</sup>2p<sup>6</sup>3s<sup>2</sup>3p<sup>6</sup>4s<sup>2</sup>3d<sup>10</sup>4p<sup>3</sup>
- ex: Strontium

# Exceptions To The Aufbau Principle

Half filled & filled subshells provide additional stability

Cr and Cu ½ fill/fill their 3d shell before the 4s shell.

Elements in same columns as Cr & Cu behave in same way.


Similar behavior seen in p block.

Cr & Cu are the only exceptions you need to know.

|                                   |         | 3d | 4s |                                      |
|-----------------------------------|---------|----|----|--------------------------------------|
|                                   | Sc [Ar] |    |    | [Ar]3d <sup>1</sup> 4s <sup>2</sup>  |
|                                   | Ti [Ar] |    |    | [Ar]3d <sup>2</sup> 4s <sup>2</sup>  |
|                                   | V [Ar]  |    |    | [Ar]3d <sup>3</sup> 4s <sup>2</sup>  |
| Cr: 3d ½ full<br>& 4s ½ full<br>→ | Cr [Ar] |    |    | [Ar]3d <sup>5</sup> 4s <sup>1</sup>  |
|                                   | Mn [Ar] |    |    | [Ar]3d <sup>5</sup> 4s <sup>2</sup>  |
|                                   | Fe [Ar] |    |    | [Ar]3d <sup>6</sup> 4s <sup>2</sup>  |
|                                   | Co [Ar] |    |    | [Ar]3d <sup>7</sup> 4s <sup>2</sup>  |
|                                   | Ni [Ar] |    |    | [Ar]3d <sup>8</sup> 4s <sup>2</sup>  |
| Cu: 3d full<br>& 4s ½ full<br>→   | Cu [Ar] |    |    | [Ar]3d <sup>10</sup> 4s <sup>1</sup> |
|                                   | Zn [Ar] |    |    | [Ar]3d <sup>10</sup> 4s <sup>2</sup> |

# Magnetism in Multi-electron Atoms


- $+1/2$  &  $-1/2$  spins will cancel if electrons paired
- No magnetic properties without spin present
  - # unpaired electrons proportional to magnetic properties



## Diamagnetic

All electrons paired

Ne:  $1s^2 2s^2 2p^6$



## Paramagnetic

At least 1 unpaired electron

Fe:  $1s^2 2s^2 2p^6 3s^2 3p^6 4s^2 3d^6$