



Chapter Fourteen: Chemical Kinetics



Rate of Reaction = speed



Kinetics = study of how fast reactions take place

Collision Theory


Three things must happen for a reaction to occur:

1. Reacting molecules must collide

Can increase the number of collisions with

- increased reactant concentration
- higher temps (= faster molecules)

2. Molecules must have the correct orientation





3. Activation Energy (E_a) must be exceeded

E_a = minimum energy required for a reaction to occur

Energy Diagrams: Activation Energy & Transition State

Given the following reaction:



Reactants: Original components before a reaction


Products: New elements/compounds formed from reaction

Transition State (aka Activated Complex): Temporary & generally unstable combination of atoms formed from molecular collision. Will be transformed into products as the reaction proceeds.

Energy Diagrams: Thermodynamics & Reaction Progress




Exothermic Reaction



Reaction progress

Net Energy Released

Endothermic Reaction




Reaction progress

Net Energy Absorbed

Reaction Rates

Kinetics:

- Study of rates of chemical reactions
- How fast does a reaction proceed?



Reaction Rate:

$$\text{Rate} = \frac{\Delta \text{Concentration}}{\Delta \text{Time}} = \frac{[\text{mol/L}]}{\Delta T} = \frac{\text{M}}{\text{sec}}$$




Rate of formation of product:

$$\text{Rate} = +\Delta[\text{O}_2]/\text{s}$$

Rate of disappearance of reactant:

$$\text{Rate} = -\Delta[\text{H}_2\text{O}_2]/\text{s}$$

(Use of negative makes rate positive, sometimes rate is written as negative)



Linearity of Reaction Rates

Reaction rates vary greatly



- Very fast; ex: burning
- Very slow; ex: disintegration of plastic in sunlight

Often not linear

- Rate is often faster at the beginning
- Rate is slowest at the end
- Some reactant often left over
- Especially pronounced for slow reactions

Average Rate of Reaction:

$$\text{Rate} = \frac{\text{Final Concentration} - \text{Initial Concentration}}{\text{Elapsed Time}}$$



Types of Reaction Rates

Reaction rate = $\Delta\text{Concentration}/\Delta\text{time} = \text{slope}$

1. Plot of original data

- [Conc.] vs time
- If curved = rate changes
 - Will reach equilibrium
- If linear = rate is constant
 - Will run out of reactant


2. Average rate of reaction

3. Instantaneous rate

- Tangent to curve 1
- Can pick any time

4. Initial rate

- $t=0$ to $t=\text{given time}$
- Generally portion of curve 1 that is linear




Instantaneous Rate



Brown color of Br_2 disappears as reaction progresses



To Find Instantaneous Rate:

- Select time
- Draw line tangent to curve at that time
- Find slope of the tangent line
- Units of [conc.]/time



Initial Rate

Portion of original data where curve is linear



Stoichiometry & Reaction Rate

When writing rates based on different reactants or products, think about what is happening



Rate at which Br_2 disappears: $X \text{ M/s}$

In terms of Br_2 :

What is the rate of formation of Br^- ? $2X \text{ M/s}$

What is the rate of disappearance of HCOOH ? $X \text{ M/s}$

Rate at which Br^- forms: $Z \text{ M/s}$

In terms of Br^- :

What is the rate of disappearance of Br_2 ? $\frac{1}{2} Z \text{ M/s}$

What is the rate of formation of CO_2 ? $\frac{1}{2} Z \text{ M/s}$

This method works for zero order & instantaneous rates, not for more complex rates.

Stoichiometry & Reaction Rate


1. The rate of decomposition of N_2O_5 at a particular instant in a reaction vessel is $4.2 \times 10^{-7} \text{ M/s}$. What is the rate of appearance of (a) NO_2 and (b) O_2 ?



(a) $8.4 \times 10^{-7} \text{ M/s}$

(b) $2.1 \times 10^{-7} \text{ M/s}$

2. Consider the reaction:




Suppose that at a particular moment during the reaction hydrogen is reacting at the rate of 0.074 M/s . (a) At what rate is ammonia being formed? (b) At what rate is nitrogen reacting?

(a) 0.049 M/s

(b) -0.025 M/s

Factors Affecting Reaction Rates

Increasing # collisions increases rate



1. Increase concentration of reactant

More particles in same volume = greater chance of collision


2. Increase temperature

Particles move faster - more likely to collide with enough E_A

3. Add a catalyst

Catalyst: chemical that does not change the amount of overall product or reactants, only the rate of the reaction

- fix molecular orientation
- alter transition state; lower E_a
- increase desired surface area



4. Improve Mixing

More interactions = greater chance of effective collision

Rate Laws:

Show Impact of Reactant Concentration on Rate

Rate Law Format: For the reaction $A + B + C \dots \rightarrow$ Products

$$\text{Rate Law: Rate} = k[A]^m[B]^n[C]^p\dots$$

Based on initial concentrations and rate

Variables required in Rate Law:

k = rate constant

m = order of reaction in reactant A

n = order of reaction in reactant B

p = order of reaction in reactant C

other letters as needed for additional reactants

Overall Order of Reaction = sum of the orders of reaction
for each reactant

$$\text{Overall Order} = m + n + p$$

Rate Laws account for non-linearity of reactions

Rate Laws con't

Properties of Rate Laws:

- Orders do not need to be positive integers
 - Ex: if concentration does not impact rate, order is zero
- Orders must be found experimentally
- Catalyst concentrations are included
- Rate Laws hold only for a specific temperature
- **Rate Laws do NOT come from the balanced equation!**

Example:



Not squared!



Rate Constant, k

- Depends on reaction, temperature, and catalyst
- Also found experimentally

Determining Rate Laws: Isolation Method

Conduct a series of experiments

- Change the concentration of just one reactant each time
- Measure the impact on the rate from each reactant
- Use the ratios of the rates to determine m, n, etc.
- Rearrange the Rate Law & plug in values to solve for k.

Example: For the reaction $A + B \rightarrow \text{Products}$ Rate = $k[A]^m[B]^n$

Conduct 3 experiments:

1. Use [A] & [B], measure Rate 1
2. Use $[A]_2$ & [B], measure Rate 2
3. Use [A] & $[B]_2$, measure Rate 3

To determine m, divide

$$\frac{\text{Rate 1}}{\text{Rate 2}} = \frac{k[A]^m[B]^n}{k[A]_2^m[B]^n} = \frac{[A]^m}{[A]_2^m}$$

To determine n, divide

$$\frac{\text{Rate 1}}{\text{Rate 3}} = \frac{k[A]^m[B]^n}{k[A]^m[B]_2^n} = \frac{[B]^n}{[B]_2^n}$$


To determine k, solve

$$k = \frac{\text{Rate}}{[A]^m[B]^n}$$

When reporting Rate Laws, use the numbers for k, m, & n

Determining Rate Laws: Isolation Method

A set of experiments was run to study the rate of the reaction:



The following data was obtained:

| Expt # | [BrO ₃ ⁻] | [Br ⁻] | [H ⁺] | Rate (M/s) |
|--------|----------------------------------|--------------------|-------------------|-------------------------|
| 1 | 0.10 | 0.10 | 0.10 | -1.2 × 10 ⁻³ |
| 2 | 0.20 | 0.10 | 0.10 | -2.4 × 10 ⁻³ |
| 3 | 0.20 | 0.30 | 0.10 | -7.4 × 10 ⁻³ |
| 4 | 0.20 | 0.10 | 0.15 | -5.4 × 10 ⁻³ |

$$\text{Rate} = k[\text{BrO}_3^-]^m[\text{Br}^-]^n[\text{H}^+]^p$$

Determining Rate Laws: Order of Reactants

$$\text{Rate} = k[\text{BrO}_3^-]^m[\text{Br}^-]^n[\text{H}^+]^p$$

| Expt # | [BrO ₃ ⁻] | [Br ⁻] | [H ⁺] | Rate (M/s) |
|--------|----------------------------------|--------------------|-------------------|-------------------------|
| 1 | 0.10 | 0.10 | 0.10 | -1.2 × 10 ⁻³ |
| 2 | 0.20 | 0.10 | 0.10 | -2.4 × 10 ⁻³ |
| 3 | 0.20 | 0.30 | 0.10 | -7.4 × 10 ⁻³ |
| 4 | 0.20 | 0.10 | 0.15 | -5.4 × 10 ⁻³ |

Order of reaction in BrO₃⁻ (m):

Order of reaction in Br⁻ (n):

Order of reaction in H⁺ (p):

Determining Rate Laws: k & Rate Law

| Expt # | [BrO ₃ ⁻] | [Br ⁻] | [H ⁺] | Rate (M/s) |
|--------|----------------------------------|--------------------|-------------------|-------------------------|
| 1 | 0.10 | 0.10 | 0.10 | -1.2 × 10 ⁻³ |
| 2 | 0.20 | 0.10 | 0.10 | -2.4 × 10 ⁻³ |
| 3 | 0.20 | 0.30 | 0.10 | -7.4 × 10 ⁻³ |
| 4 | 0.20 | 0.10 | 0.15 | -5.4 × 10 ⁻³ |

$$\text{Rate} = k[\text{BrO}_3^-]^m[\text{Br}^-]^n[\text{H}^+]^p$$

$$m = 1$$

$$n = 1$$

$$p = 2$$

k: Solve using data from each experiment (4X!) & average

$$k = \frac{\text{Rate}}{[\text{BrO}_3^-]^1[\text{Br}^-]^1[\text{H}^+]^2}$$

Rate Law =

Rate Laws can be used to determine the initial speed of a reaction with given concentrations of reactants.

For the reaction:



we determined that the Rate Law was:

$$\text{Rate} = -12\text{M}^{-3}\text{s}^{-1}[\text{BrO}_3^-] [\text{Br}^-] [\text{H}^+]^2$$

If we start with the following concentrations:

$$[\text{BrO}_3^-] = 0.4\text{M}$$

$$[\text{Br}^-] = 0.8\text{M}$$

$$[\text{H}^+] = 0.2\text{M}$$

How fast will the initial reaction proceed?

$$\begin{aligned}\text{Rate} &= -12\text{M}^{-3}\text{s}^{-1} \times [0.4\text{M}] \times [0.8\text{M}] \times [0.2\text{M}]^2 \\ &= 0.15 \text{M}^{-3}\text{s}^{-1} \text{M}^4 \\ &= 0.15 \text{Ms}^{-1}\end{aligned}$$

Types of Reactions Based on Rate: Zero Order

Change in concentration over time IS linear

Rate Law for $A \rightarrow$ Product: $\text{rate} = k [A]^0 = k$


Units of k : $k = \text{rate} = \text{M/s}$

Linear Rate Equation:

$$[A] = -kt + [A]_0$$

Eventually $[A]$ will go to zero
(all reactant will be used up)

Note that this equation is in the form $y = mx + b$, where b is the y -intercept (i.e. the initial concentration of A).



Types of Reactions Based on Rate: First Order

Change in concentration over time is NOT linear

Change in the natural log (ln) of concentration over time IS linear.


Rate Law for $A \rightarrow$ Product: $\text{rate} = -\frac{\Delta[A]}{\Delta t} = k[A]$

Units of k : $k = \text{rate}/[A] = (\text{M/s})/\text{M} = 1/\text{s} = \text{s}^{-1}$

$[A]$ vs. time is Nonlinear

$\ln [A]$ vs time is Linear:


$$[A] = [A]_0 e^{(-kt)}$$



$$\ln \left(\frac{[A]}{[A]_0} \right) = -kt$$

$$[A]_t = [A] \text{ at time } t$$
$$[A]_0 = [A] \text{ at } t=0$$

$$\ln[A] = -kt + \ln[A]_0$$



Types of Reactions Based on Rate: Second Order

Change in concentration over time is NOT linear


Change in the inverse of concentration ($1/[A]$) over time IS linear.

Rate Law for $A \rightarrow$ Product: $\text{rate} = -\frac{\Delta[A]}{\Delta t} = k[A]^2$

Units of k : $k = \text{rate}/[A]^2 = (\text{M/s})/\text{M}^2 = 1/\text{Ms} = \text{M}^{-1}\text{s}^{-1}$

The linear equation is:

$$\frac{1}{[A]} = kt + \frac{1}{[A]_0}$$



Predicting Reaction Order Graphically


For the reaction $A \rightarrow \text{Products}$:

1. Calculate $\ln[A]$ and $1/[A]$
2. Make 3 graphs: $[A]$, $\ln[A]$, and $1/[A]$ vs. time
3. The graph with the best straight line is the reaction order

| | Zero | First | Second |
|---------|---------|-----------|---------|
| time, s | $[A]$ M | $\ln [A]$ | $1/[A]$ |
| 0 | 1.710 | 0.536 | 0.585 |
| 4 | 1.150 | 0.140 | 0.870 |
| 8 | 0.870 | -0.139 | 1.149 |
| 24 | 0.460 | -0.777 | 2.174 |
| 45 | 0.280 | -1.273 | 3.571 |

Predicting Reaction Order Graphically


[A] vs. time



$$[A] = -kt + [A]_0$$

Not linear, so not Zero Order


ln[A] vs. time



$$\ln[A] = -kt + \ln[A]_0$$

Not linear, so not First Order

1/[A] vs. time



$$1/[A] = kt + 1/[A]_0$$

Linear, so reaction is


Second Order

Using a Graph to Find Rate Constant, k

Equation of the line is: $y = mx + b$

$$1/[A] = kt + 1/[A]_0$$

$k = m = \text{slope!}$



$$k = m = (y - y_1)/(x - x_1)$$

$$= (3.57 - 1.15)/(45 - 8)$$

$$= 0.066 \text{ M}^{-1}\text{s}^{-1}$$

$$\frac{1}{[A]} = 0.066t + \frac{1}{[A]_0}$$

Using Line Equations to Find Information

1. For the first order reaction: $2 \text{N}_2\text{O}_5 \rightarrow 2 \text{N}_2\text{O}_4 + \text{O}_2$
at 45°C , $k = 6.22 \times 10^{-4} \text{ s}^{-1}$. If $[\text{N}_2\text{O}_5] = 0.100\text{M}$:

a.) How long does it take for the concentration to drop to 0.010M ? A: $3.70 \times 10^3\text{s}$

b.) What is the concentration after one hour? A: 0.0107M

2. The decomposition of sulfuryl chloride (SO_2Cl_2) is first order in SO_2Cl_2 . The rate constant for the decomposition at 660K is $4.5 \times 10^{-2} \text{ s}^{-1}$.

a.) If we begin with an initial SO_2Cl_2 pressure of 450 torr, what is the pressure of this substance after 60.s? **A: 30. torr**


b.) At what time will the pressure of SO_2Cl_2 decline to 1/10 its initial value? **A: 51s**

Half-Life ($t_{1/2}$)

The time required for $[A]_0$ to decrease by 50%

- NOT the time for half of the reaction to occur
- Provides another way to get the rate constant

- Symbol is $t_{1/2}$
- $t_{1/2}$ depends on reaction order
- Can solve for $t_{1/2}$ by plugging $[A]_0 = 1A$ and $[A] = \frac{1}{2}A$ into rate equation



Half-Life of a First Order Reaction

Linear Equation:

$$\ln[A] = -kt + \ln[A]_0$$

Rearranged Equation:


$$\ln \frac{[A]}{[A]_0} = -kt$$

Plug in: $[A]_0 = 1A$ & $[A] = \frac{1}{2}A$

$$\ln(\frac{1}{2}A/A) = \ln(1/2) = -kt_{1/2}$$

Solve for $t_{1/2}$:

$$t_{1/2} = \ln 2 / k = \mathbf{0.693 / k}$$




For a First Order Reaction:

$t_{1/2}$ is independent of initial reactant concentration

If you know $t_{1/2}$, you know the rate constant

Summary of Kinetic Data

| Order | Rate Law | Integrated Rate Law | Linear Plot | k | Half-life |
|-------|--------------------------|-----------------------------|----------------|--------|------------|
| 0 | Rate = k | $[A]_t = -kt + [A]_0$ | [A] vs. t | -slope | $[A]_0/2k$ |
| 1 | Rate = k[A] | $\ln[A]_t = -kt + \ln[A]_0$ | $\ln[A]$ vs. t | -slope | $0.693/k$ |
| 2 | Rate = k[A] ² | $1/[A]_t = kt + 1/[A]_0$ | $1/[A]$ vs. t | +slope | $1/k[A]_0$ |



Half-life Examples

1. For a 1st order reaction, determine the time it will take for only 1/8 of a material to be left if $t_{1/2} = 3.47 \times 10^2 \text{s}$

A: $1.04 \times 10^3 \text{s}$


2. The reaction below is first order in $[\text{H}_2\text{O}_2]$:



A solution originally at 0.600M is found to be 0.075M after 54min. Determine the half-life.

A: 18 min

3. The thermal decomposition of phosphine (PH₃) into phosphorus and molecular hydrogen is a first order reaction:



The half-life of the reaction is 35.0s at 680°C. Calculate (a) the first order rate constant for the reaction and (b) the time required for 95% of the phosphine to decompose.

A: (a) 0.0198s⁻¹ (b) 151s

4. The rate constant for the second-order reaction



is $0.54 \text{ M}^{-1}\text{s}^{-1}$ at 300°C .

(a) How long, in seconds, would it take for the concentration of NO_2 to decrease from 0.62M to 0.28M ?

A: 3.6s

(b) Calculate the half-lives at these two concentrations.

A: 3.0s, 6.6s

Effect of Temperature on Reaction Rate: Arrhenius Equation

Arrhenius Equation – Shows how temp. affects the rate constant (k)

$$k = Ae^{-E_a/RT} \quad \text{or} \quad \ln k = (-E_a/RT) + \ln A$$

k = rate constant

A = frequency factor (based on collision frequency & the probability of correct orientation; constant for a given system)

E_a = activation energy

R = gas constant in energy units ($8.314 \text{ J mol}^{-1} \text{ K}^{-1}$)

T = temperature in Kelvin

Graphing **$\ln k$ vs. $1/T$** gives a straight line with
slope = $-E_a/R$


Provides a method to determine E_a for a reaction

Arrhenius Equation Example



Step 1: Find k at several temps. Step 2: Graph $\ln k$ vs. $1/T$

| Temp. (K) | k ($\text{M}^{-1}\text{s}^{-1}$) |
|-----------|--------------------------------------|
| 283 | 3.52×10^{-7} |
| 356 | 3.02×10^{-5} |
| 393 | 2.19×10^{-4} |
| 427 | 1.16×10^{-3} |
| 508 | 3.95×10^{-2} |



Step 3: Calculate slope ($= -E_a/R$) Step 4: Multiply slope by $-R$ to get E_a

2 -Point Arrhenius Equation Example

Use rate constants at just 2 temps to solve for E_a

- This method is less accurate but only requires 2 k values
- Best to use **only if there is limited data**

- Equation:
$$\ln \frac{k_1}{k_2} = \frac{E_a}{R} \left(\frac{1}{T_2} - \frac{1}{T_1} \right)$$
 Note:
 $\ln(k_1/k_2) = \ln k_1 - \ln k_2$

Find E_a for the reaction: $2\text{HI}(\text{g}) \rightarrow \text{H}_2(\text{g}) + \text{I}_2(\text{g})$

Data:

$$k_1 = 0.0395 \text{ M}^{-1}\text{s}^{-1}$$

$$T_1 = 508 \text{ K}$$

$$k_2 = 0.00116 \text{ M}^{-1}\text{s}^{-1}$$

$$T_2 = 427 \text{ K}$$

$E_a = 78.5 \text{ kJ/mol}$ – note difference from previous slide!

More uses of the Arrhenius Equation

1. The activation energy of a certain reaction is 31.5 kJ/mol. At 30°C, the rate constant is 0.0190 s⁻¹. At what temperature in degrees Celsius would this reaction go twice as fast?

A: 48°C

2. The gas-phase reaction $\text{Cl(g)} + \text{HBr(g)} \rightarrow \text{HCl(g)} + \text{Br(g)}$ has an overall enthalpy change of -66kJ . The activation energy for the reaction is 7 kJ .



(a) Sketch the energy profile for the reaction, and label E_a and ΔE .

(b) What is the activation energy for the reverse reaction?

A: 73 kJ

3. On the basis of the frequency factors and activation energy values of the following two reactions, determine which one will have the larger rate constant at room temperature (298K).



A: second reaction




Reaction Mechanism: What (we think) actually happens during a reaction

Series of elementary reactions that lead to an overall rxn

Mechanism info can be used to modify reactions to increase yield, speed, etc.




N_2O_2 is detected during the reaction!



Some Terminology: **Molecularity**

Number of molecules reacting in an elementary step


Unimolecular



Unimolecular – one

- Fastest
- Molecule simply breaks apart
- No molecular alignment needed


Bimolecular



Bimolecular – two

- Most common
- May require correct alignment

Termolecular



Termolecular – three

- Usually rate limiting (slow!)
- Low probability that all 3 particles will collide simultaneously (esp. with correct alignment and energy)

Some More Terminology

Intermediates: Species that appear in a reaction mechanism but not in the overall reaction


- Formed in an early elementary step
- Consumed in a later elementary step

Ex:


Elementary Step:



+ Elementary Step:



Overall Reaction:



N_2O_2 is an intermediate

Rate Determining Step: Slowest step in the sequence of steps leading to the overall reaction

- Often determined by molecularity

Rate Laws & Intermediate Steps

Unimolecular reaction: $A \rightarrow \text{Products}$ rate = $k[A]$


Bimolecular reaction: $A + B \rightarrow \text{Products}$ rate = $k[A][B]$

Bimolecular reaction: $A + A \rightarrow \text{Products}$ rate = $k[A]^2$

Writing reaction mechanisms:

- The sum of the elementary reactions must give the overall balanced equation for the reaction.
- The rate-determining step should predict the same rate law that is determined experimentally
- The rate law for each step is found from the coefficients of the reactants. This is **ONLY** true for **ELEMENTARY** reactions!

Developing a Reaction Mechanism



Measure the rate of the reaction:

- Run a set of isolation method experiments
 - Run 1 experiment for each reactant + a control
 - Measure $[A]$ vs. time

Formulate the rate law

- Determine the order of each reactant based on data
- Use the results to calculate an average value for k
- Determine the overall rate of the reaction

Suggest a reaction mechanism

- Elementary steps must match stoichiometry of overall rxn
- Use molecularity to determine rate determining step
- Order of rate-determining step must match overall rxn order

Ex: Developing a Reaction Mechanism for


$$\text{NO}_2 + \text{CO} \rightarrow \text{NO} + \text{CO}_2$$

Based on the results of a set of isolation method experiments, the overall rate was found to be:

$$\text{Rate} = k[\text{NO}_2]^2$$

Suggested reaction mechanism:

- Need to account for rate being 2nd order in NO₂
- Need to account for rate being zero order in CO



Step 1 would be the rate-determining step, because the rate matches the overall reaction rate.






You might also be asked to evaluate possible mechanisms.

Evaluating Mechanisms







Rate = $k[\text{H}_2][\text{ICl}]$

Option #1







Rate = $k[\text{ICl}]^2$ ✗

Option #3






Rate = $k[\text{H}_2]$ ✗

Option #2



Rate = $k[\text{H}_2][\text{ICl}]$ ✓

Option #4



Rate = $k[\text{H}_2][\text{ICl}]$ ✓

Catalysis

Catalysts alter the rate of a chemical reaction

- Not part of the overall reaction (not a reactant or product)
- Often work by lowering activation energy
 - Since less energy is needed, rate increases
- Materials that slow reaction rates are called **inhibitors**


Heterogeneous catalysts

- In a different phase than the reacting medium
- Solids often used to catalyze reactants in solution
- Often used as a surface to align molecules correctly
- Benefit – often easy to remove & re-use


Homogeneous catalysts

- In the same phase as the reacting medium
- May have larger impact on rate (may be able to create a lower energy activated complex than heterogeneous)

Effect of a Catalyst on Reaction Profile & Activation Energy




Effect of a Catalyst on Reaction Profile & Activation Energy



Enzymes

Enzymes

- Control the rate of biological reactions
- Very specific
 - Only act on specific molecules called **substrates**
- Often proteins – shape determines functionality
 - Partially due to hydrogen bonding (CHM101!)
- Some enzymes also act as **inhibitors**



Catalysts & Reaction Mechanisms

When considering mechanisms:

- Catalysts control the rate of the reaction.
- The rate of the elementary step that involves the catalyst will control the observed rate.
- Rate constant is different from the rate constant without the catalyst - called the catalytic rate constant.

