


# Chapter 16

## Chemical Equilibrium




# Equilibrium: A Dynamic Process

## Opposing processes occur at equal rates

- Forward & reverse reactions occur at equal rates
- No outward change is observed
- **Ratio** of reactants to products is constant
- Temperature dependent
- Other factors can also shift equilibrium toward products or reactants
- Represented by double arrows ( $\rightleftharpoons$  or  $\longleftrightarrow$ )


## Physical Equilibrium

Ex: Equilibrium between phases



## Chemical Equilibrium


Equilibrium between reactants & products



Initial conditions may vary – concentrations will adjust to establish equilibrium




As a system **approaches** equilibrium, both the forward and reverse reactions are occurring



At **equilibrium**, the forward and reverse reactions are proceeding at the same rate, so the relative concentrations remain constant. <sub>3</sub>

# Equilibrium Constant ( $K_c$ )



At equilibrium,  $[\text{N}_2\text{O}_4]$  &  $[\text{NO}_2]$  are constant

- **NOT EQUAL**
- Not static – both forward & reverse reactions continue
- Actual amounts depend on system

Rate (forward) = Rate (reverse):  $k_1[\text{N}_2\text{O}_4] = k_{-1}[\text{NO}_2]^2$

$$\frac{k_1}{k_{-1}} = \frac{[\text{NO}_2]^2}{[\text{N}_2\text{O}_4]} = K_c$$

**TABLE 15.1** The  $\text{NO}_2$ – $\text{N}_2\text{O}_4$  System at 25° C

Initial Concentrations (M)		Equilibrium Concentrations (M)		Ratio of Concentrations at Equilibrium	
$[\text{NO}_2]$	$[\text{N}_2\text{O}_4]$	$[\text{NO}_2]$	$[\text{N}_2\text{O}_4]$	$\frac{[\text{NO}_2]}{[\text{N}_2\text{O}_4]}$	$\frac{[\text{NO}_2]^2}{[\text{N}_2\text{O}_4]}$
0.000	0.670	0.0547	0.643	0.0851	$4.65 \times 10^{-3}$
0.0500	0.446	0.0457	0.448	0.102	$4.66 \times 10^{-3}$
0.0300	0.500	0.0475	0.491	0.0967	$4.60 \times 10^{-3}$
0.0400	0.600	0.0523	0.594	0.0880	$4.60 \times 10^{-3}$
0.200	0.000	0.0204	0.0898	0.227	$4.63 \times 10^{-3}$

[conc]  
not equal

Ratio  
is equal

# Equilibrium Expression

For the reaction:  $aA + bB + \dots \rightleftharpoons cC + dD + \dots$

The Equilibrium Expression is: 
$$K_c = \frac{[C]^c[D]^d}{[A]^a[B]^b}$$

For an Equilibrium Expression:

- Concentrations of products are in numerator
- Concentrations of reactants are in denominator
- They are the concentrations at equilibrium
- Exponents ARE coefficients from balanced equation
- Units generally not included
- Also known as a Mass Action Expression

Note difference from rate equation – equilibrium expression IS BASED ON BALANCED EQUATION

# Impact of How an Equation is Balanced

Reaction A



$$K_c = \frac{[\text{SO}_3]}{[\text{SO}_2][\text{O}_2]^{1/2}}$$

Reaction B



$$K_c = \frac{[\text{SO}_3]^2}{[\text{SO}_2]^2[\text{O}_2]}$$

Equilibrium constants change if the reaction is balanced differently

Numerical values for  $K_c$  are related, but different

$$K_c (\text{reaction B}) = [K_c (\text{reaction A})]^2$$

It is essential to know how the reaction was balanced

# $K_p$ The Pressure Version of $K_c$

Remember the Gas Laws – Chapter 5!



- In a closed system, pressure  $\propto$  concentration
  - The equilibrium expression can also be written in terms of pressure
  - Very useful since gas phase reactions are often monitored via pressure

$$K_p = \frac{(P_C)^c (P_D)^d}{(P_A)^a (P_B)^b}$$

$\propto$  = proportional to

# Writing Equilibrium Expressions

Write the equilibrium constant expression  $K_p$  and  $K_c$  for:






# Manipulating Chemical Equations & $K_c$


When reversing a chemical equation, invert  $K_c$



When multiplying coefficients by  $n$ , raise  $K_c$  to  $n^{\text{th}}$  power




When adding equations, multiply the  $K_c$  values



# Manipulating K values

1. For the formation of  $\text{NH}_3$  from  $\text{N}_2$  and  $\text{H}_2$



$K_p = 4.34 \times 10^{-3}$  at  $300^\circ\text{C}$ . What is the value of  $K_p$  for the reverse reaction?

A:  $2.30 \times 10^2$

2. How does the magnitude of the equilibrium constant  $K_p$  for the reaction



change if the equilibrium is written as



A:  $K_p$  is cubed

3. Given that, at 700 K,  $K_p = 54.0$  for the reaction:



and  $K_p = 1.04 \times 10^{-4}$  for the reaction:



determine the value of  $K_p$  for the following reaction at 700K:




# Types of Equilibria: Homogeneous ( $K_c$ & $K_p$ )

All products & reactants are in the same phase

$K_c$ : All chemicals are in units of molarity (mol/L)  
Can be either **gas** or **aqueous** solutions

$K_p$ : All chemicals in **gas** phase  
Pressure is the partial pressure  
- units often atmospheres



$K_c$  &  $K_p$  are related:  $K_p = K_c \cdot (RT)^{\Delta n}$

- Derived from ideal gas law
- $R$  = gas constant (0.0821 L · atm/mol · K)
- $T$  = temperature in Kelvin
- $\Delta n$  = moles gaseous products - moles gaseous reactants

$K_p = K_c$  if moles product gas = moles reactant gas

If moles product gas  $\neq$  moles reactant gas, there will be a change in pressure

## Calculating $K_p$ from $K_c$

Calculate  $K_p$  for the following reactions at  $1500^\circ\text{C}$ .



$$K_c = 12800$$

$$\Delta n_{\text{gas}} = -1$$



$$K_c = 12800$$

$$\Delta n_{\text{gas}} = 0$$

$$K_p = K_c \cdot (RT)^{\Delta n}$$

A: 87.9

A: 12800

# Types of Equilibria: Heterogeneous

Are the equilibrium constants for the following two reactions the same?



$$“K_1” = \frac{[\text{H}_2\text{SO}_4(\text{g})]}{[\text{SO}_3(\text{g})] [\text{H}_2\text{O}(\text{g})]}$$

$$“K_2” = \frac{[\text{H}_2\text{SO}_4(\text{aq})]}{[\text{SO}_3(\text{g})] [\text{H}_2\text{O}(\text{l})]}$$


“K<sub>1</sub>” would equal “K<sub>2</sub>” ONLY IF [H<sub>2</sub>SO<sub>4</sub>(g)] = [H<sub>2</sub>SO<sub>4</sub>(aq)]  
AND [H<sub>2</sub>O(g)] = [H<sub>2</sub>O(l)]

[H <sub>2</sub> SO <sub>4</sub> (g)]	dependent on partial pressure	variable
[H <sub>2</sub> SO <sub>4</sub> (aq)]	# moles dissolved in water	variable
[H <sub>2</sub> O(g)]	dependent on partial pressure	variable
[H <sub>2</sub> O(l)]	1 g/mL = 1000 g/L	56M
	for 1L: 1000g(1 mol/18g) = 56 mol	

**The two equilibrium constants are NOT the same!**

**THE PHASE MATTERS!**

# Equilibria Involving Pure Solids & Liquids




$P_{\text{CO}_2}$  does not depend on the amount of  $\text{CaCO}_3$  or  $\text{CaO}$   
As long as there is some  $\text{CaCO}_3$  &  $\text{CaO}$  in the system, the amount of  $\text{CO}_2$  above the solid will remain the same

# Equilibria Involving Mixed Phases (i.e. Heterogeneous)

The subscript “eq” is for general equilibria

$K_c$  &  $K_p$  are specific types of  $K_{eq}$

Equilibrium expressions only include gaseous & aqueous materials (solids & liquid are NOT included)



$K_p$  for liquid-vapor equilibrium = vapor pressure of liquid




To write an appropriate equilibrium expression from a balanced equation, the equation **MUST** show the phase of each reactant & product.



## Determination of $K_{\text{eq}}$ values:

- The equilibrium expression comes from the balanced equation



- The value of  $K$  is experimentally determined
  - Allow the reaction to run until it reaches equilibrium.
  - Once equilibrium is reached, measure the concentration of all components in the equilibrium expression.
  - Plug the equilibrium concentrations into the equilibrium expression to calculate  $K_{\text{eq}}$ .

# Equilibria Involving Solids & Liquids

A mixture of  $\text{H}_2$ ,  $\text{S}$ , and  $\text{H}_2\text{S}$  is held in a 1.0L vessel at  $90^\circ\text{C}$  and reacts according to the equation:



An equilibrium mixture contains 0.46g  $\text{H}_2\text{S}$  & 0.40g  $\text{H}_2$ .


(a) Write the equilibrium constant expression for this reaction.

(b) What is the value of  $K_c$  for the reaction at this temp.?

A: 0.068

(c) Why can we ignore the amount of  $\text{S}$  when doing the calculation in part b?



# What Equilibrium Constants Can Tell Us





At Equilibrium:

$$K_c = \frac{[C]^c [D]^d}{[A]^a [B]^b}$$

$K > 1$   
More Products  
than Reactants



$K < 1$   
More Reactants  
than Products



As  $K$  goes to infinity, reaction goes to completion.  
As  $K$  goes to zero, no reaction occurs.

# Analyzing $K_p/K_c$ Values

For the reaction:  $\text{H}_2(\text{g}) + \text{I}_2(\text{g}) \rightleftharpoons 2 \text{HI}(\text{g})$

$K_p = 794$  at 298K and  $K_p = 54$  at 700K

Is the formation of HI more favored at the higher or lower temperature?

# Using the Equilibrium Expression

1. Nitrogen monoxide exists in equilibrium with nitrogen and oxygen gas. At a given temperature, 0.100 moles of NO were added to a 2.00L vessel. At equilibrium, 0.044 moles of NO were remaining. What is the value of  $K_c$ ?


1. Write the balanced equation & equilibrium constant expression.

2. Find equilibrium molarities of reactants and products.

We are using concentration, so make sure to convert to Molarity!

3. Calculate  $K_c$

# Predicting the Direction of a Reaction



## If the value of $K_{eq}$ is very large


- [C] & [D] very large or [A] & [B] very small
- Reaction will go to completion
- Equilibrium favors products

## If the value of $K_{eq}$ is very small

- [C] & [D] very small or [A] & [B] very large
- Reaction proceeds very slowly or no reaction (NR) at all
- Equilibrium favors reactants

Note that  $K_{eq}$  has its origins in thermodynamics, but kinetics are also important. The value of  $K_{eq}$  does not guarantee that a reaction will occur in either direction.

# The Reaction Quotient ( $Q_c$ or $Q_p$ )




Essentially  $K_{eq}$  but at non-equilibrium conditions

- Indicates direction of reaction to get to equilibrium


$Q < K$

- $[Products] <$  equilibrium concentration
- Higher rate in forward direction




$Q > K$

- $[Products] >$  equilibrium concentration
- Higher rate in reverse direction




$Q = K$

- $[Products] =$  equilibrium concentration
- Forward rate = Reverse rate
- No macroscopic changes observed



# Reaction Quotient

1. For the reaction of hydrogen with nitrogen to form ammonia at 25°C,  $K_c = 4.0 \times 10^8$ . The equilibrium concentrations were found to be  $1.0 \times 10^{-3} \text{M}$  in both hydrogen & nitrogen and  $0.020 \text{M}$  in ammonia. If you add  $0.010 \text{M}$  nitrogen, in which direction does the reaction shift?





2. At 1000K, the value of  $K_p$  for the following reaction is 0.338.



Given the following pressures:

$$P_{\text{SO}_3} = 0.16 \text{ atm}; P_{\text{SO}_2} = 0.41 \text{ atm}; P_{\text{O}_2} = 2.5 \text{ atm}$$

(a) Is the system at equilibrium? **A: No**

(b) If not, predict the direction in which the reaction will shift to reach equilibrium. **A: toward reactants**

# Free Energy & Equilibrium

During a reaction, not all of the reactants and products will be in their standard state. Under any conditions (standard or nonstandard), the free energy change is:

$$\Delta G = \Delta G^\circ + RT \ln Q$$

$$R = 8.314 \times 10^{-3} \text{ kJ/mol K}$$

T = temperature in K

Q = reaction quotient

(Under standard conditions, all concentrations are 1 M, so  $Q = 1$ ,  $\ln Q = 0$ , and  $\Delta G = \Delta G^\circ$ )

At equilibrium:  $\Delta G = 0$  and  $Q = K$  (K is equilibrium constant)

So the equation becomes:  $0 = \Delta G^\circ + RT \ln K$

Therefore, at equilibrium:  $\Delta G^\circ = -RT \ln K$

# Gibbs Free Energy & Equilibrium

$K$	$\ln K$	$\Delta G^\circ$	Comments
$> 1$	Positive	Negative	Products are favored over reactants at equilibrium.
$= 1$	0	0	Products and reactants are equally favored at equilibrium.
$< 1$	Negative	Positive	Reactants are favored over products at equilibrium.

$\Delta G^\circ < 0$  (negative), the forward reaction is spontaneous

$\Delta G^\circ > 0$  (positive), the reverse reaction is spontaneous

$\Delta G^\circ = 0$  the reaction is at equilibrium

The magnitude of  $\Delta G^\circ$  indicates how far the standard state is from equilibrium  
(more negative  $\Delta G^\circ$  = larger  $K$ )

# Free Energy & Equilibrium

1. Calculate  $\Delta G^\circ$  for the following reaction at 25°C.



$K_{\text{sp}}$  for  $\text{Fe(OH)}_2$  is  $1.6 \times 10^{-14}$

A: 79kJ/mol

2. For the following reaction:



A: (a) -5.41 kJ/mol  
(b) 0.295 kJ/mol

- (a) Using data from Appendix 2, calculate  $\Delta G^\circ$  at 298K  
(b) Calculate  $\Delta G$  at 298K if the partial pressure of  $\text{NO}_2$  and  $\text{N}_2\text{O}_4$  are 0.40atm and 1.60atm, respectively.

# Using $K_{eq}$ to Find Equilibrium Concentrations

1. Write the properly balanced chemical equation.
  - Need the stoichiometry & species involved in the reaction.
2. Set up a table of concentrations for all components.
  - Use coefficients from balanced equation to express all unknown concentrations in terms of a single reactant or product,  $x$ .
3. Write out the equilibrium constant expression for the reaction (i.e. the equation for  $K$ ).
4. Substitute the values from step 2 into the  $K_{eq}$  equation.
5. Solve the equation for  $x$ .
6. Substitute the value you calculated for  $x$  into the expressions for the other equilibrium concentrations.

**“Ice Tables” will be used in most of the remaining chapters!**

# Using $K_{eq}$ to Find Equilibrium Concentrations

1. The decomposition of BrCl to bromine & chlorine has a  $K_c$  of 0.14 at 350K. If the initial concentration of BrCl is 0.062M, what are the equilibrium concentrations of all components?

1. Balanced Equation:

2. ICE Table

3.  $K_{eq}$  Expression

6. Use  $x$  to determine conc.

4 & 5. Substitute values from table & Solve for  $x$

2. Carbon monoxide reacts with water at  $1000^{\circ}\text{C}$  to give carbon dioxide and hydrogen with  $K_c = 0.58$ . A reaction was started with the following composition:  $\text{CO}_2$  0.20M;  $\text{H}_2$  1.20M;  $\text{H}_2\text{O}$  0.50M;  $\text{CO}$  1.00M. What are the equilibrium concentrations of all components?

1. Balanced Equation:

2. ICE Table

3.  $K_{\text{eq}}$  Expression

4. Substitute values from table

Need quadratic equation!



5. Solve for x using quadratic equation:  $\frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$

6. Use x to determine concentrations

3. Compare the molar solubility of lead chloride ( $[\text{Pb}^{2+}]$ ) at 25°C and 90°C.



$K_{\text{sp}}$  for  $\text{PbCl}_2$  is  $1.6 \times 10^{-5}$  at 25°C

From Table:

$\Delta H^{\circ}_{\text{f}}$  (kJ/mol)

$S^{\circ}$  (J/molK)

$\text{PbCl}_2(\text{s}) = -359$

$\text{PbCl}_2(\text{s}) = 136$

$\text{Pb}^{2+}(\text{aq}) = -1.7$

$\text{Pb}^{2+}(\text{aq}) = 10.5$

$\text{Cl}^{-}(\text{aq}) = -167.2$

$\text{Cl}^{-}(\text{aq}) = 56.5$

Molar solubility = amt. moles dissolved = equilibrium concentration!

At 25°C, use K to solve for X

A: solubility at 90°C about double the solubility at 25°C

3. Compare the molar solubility of lead chloride ( $[\text{Pb}^{2+}]$ ) at 25°C and 90°C.



$K_{\text{sp}}$  for  $\text{PbCl}_2$  is  $1.6 \times 10^{-5}$  at 25°C

From Table:

$\Delta H^{\circ}_f$ (kJ/mol)	$S^{\circ}$ (J/molK)
$\text{PbCl}_2(\text{s}) = -359$	$\text{PbCl}_2(\text{s}) = 136$
$\text{Pb}^{2+}(\text{aq}) = -1.7$	$\text{Pb}^{2+}(\text{aq}) = 10.5$
$\text{Cl}^{-}(\text{aq}) = -167.2$	$\text{Cl}^{-}(\text{aq}) = 56.5$

Molar solubility = amt. moles dissolved = equilibrium concentration!

$$\Delta H_{\text{rxn}} = [-1.7 \text{ kJ/mol} + 2(-167.2 \text{ kJ/mol})] - [359 \text{ kJ/mol}] = 22.9 \text{ kJ/mol}$$

$$\Delta S_{\text{rxn}} = [10.5 \text{ J/molK} + 2(56.5 \text{ J/molK})] - [136 \text{ J/molK}] = -12.5 \text{ J/molK}$$

$$= -0.0125 \text{ kJ/molK}$$

At 90°C, use  $\Delta G$  to solve for K, then use the new K to solve for X

A: solubility at 90°C about double the solubility at 25°C

# Making Approximations in Equilibrium Calculations (do we need to use the Quadratic Equation?)

**Answer: Only if the percent difference between X and the equilibrium concentration is >5%**

## Example A:

Find [HClO] for a 0.010M solution of hypochlorous acid (HClO) at 25°C.  $\text{HClO(aq)} + \text{H}_2\text{O(l)} \rightleftharpoons \text{H}_3\text{O}^+(\text{aq}) + \text{ClO}^-(\text{aq})$  where  $K_a = 2.9 \times 10^{-8}$ .

	HClO	H <sub>3</sub> O <sup>+</sup>	ClO <sup>-</sup>
I	0.010	0	0
C	-X	+X	+X
E	0.010-X	X	X

$$K_a = \frac{[\text{H}_3\text{O}^+][\text{A}^-]}{[\text{HA}]}$$

$$2.9 \times 10^{-8} = \frac{[\text{X}][\text{X}]}{[0.010 - \text{X}]}$$

Ignore this X

$$2.9 \times 10^{-8} = \frac{X^2}{0.010}$$

$$X^2 = 2.9 \times 10^{-10}$$

$$X = 0.000017$$

$$[\text{HClO}] = 0.010 - 0.000017 \sim 0.010$$

$$(0.000017/0.010) * 100 = 0.17\%$$

Approximation OK

# Making Approximations in Equilibrium Calculations (do we need to use the Quadratic Equation?)

**Answer: Only if the percent difference between X and the equilibrium concentration is >5%**

## Example B:

Find  $[\text{HNO}_2]$  for a 0.010M solution of nitrous acid ( $\text{HNO}_2$ ) at  $25^\circ\text{C}$ .  $\text{HNO}_2(\text{aq}) + \text{H}_2\text{O}(\text{l}) \rightleftharpoons \text{H}_3\text{O}^+(\text{aq}) + \text{NO}_2^-(\text{aq})$  where  $K_a = 4.5 \times 10^{-4}$ .

	$\text{HNO}_2$	$\text{H}_3\text{O}^+$	$\text{NO}_2^-$
I	0.010	0	0
C	-X	+X	+X
E	$0.010 - X$	X	X

$$4.5 \times 10^{-4} = \frac{X^2}{0.010}$$

$$X^2 = 4.5 \times 10^{-6}$$

$$X = 0.0021$$

$$K_a = \frac{[\text{H}_3\text{O}^+][\text{A}^-]}{[\text{HA}]}$$

$$4.5 \times 10^{-4} = \frac{[\text{X}][\text{X}]}{[0.010 - \text{X}]}$$


Ignore this X

$$[\text{HNO}_2] = 0.010 - 0.0021 \sim 0.008$$

$$(0.0021/0.008) * 100 = 26\%$$

Approximation **NOT** acceptable

Hydrofluoric acid dissociates in water to yield hydronium ions and fluoride ions according to the equation:



The  $K_{\text{eq}}$  for this process is  $7.1 \times 10^{-4}$ . If you start with a 1.2M solution of HF, what are the equilibrium concentrations for all components?

1. Balanced Equation:




2. ICE Table

3.  $K_{\text{eq}}$  Expression

4. Substitute values from table




# Factors that Affect Chemical Equilibrium: Le Châtelier's Principle

When stress is applied to a system at equilibrium, the system will shift to reduce the applied stress and re-establish equilibrium



Addition of pressure causes the reaction to shift towards products – fewer moles of gas decreases the pressure (i.e. the stress)

# Remember: Equilibrium Can Be Reached From Either Direction



**It doesn't matter whether you start with  $\text{N}_2$  and  $\text{H}_2$  or with  $\text{NH}_3$  - you will have the same proportions of all three substances at equilibrium**



# Many different types of stress can affect equilibrium, such as:

Change in components

Change in concentrations


Change in partial pressure

Change in external pressure

Change in total volume

Change in temperature

# Changing Components, Concentration, or Partial Pressure



$$K_{eq} = \frac{[\text{Asp.}]}{[\text{AA}][\text{SA}]}$$

Synthesis of Aspirin from Salicylic Acid & Acetic Acid is an equilibrium process. We want to make as much aspirin as possible.

## Addition of a component: Add Acetic Acid


$K_{eq}$  must remain constant

$$K_{eq} = \frac{[\text{Asp.}]}{[\text{AA}][\text{SA}]}$$

If we add more AA, the denominator gets larger

To keep  $K_{eq}$  constant, we need to increase the numerator – reaction will shift toward the products (make more aspirin).

# Changing Components, Concentration, or Partial Pressure



$$K_{eq} = \frac{[\text{Asp.}]}{[\text{AA}][\text{SA}]}$$

Synthesis of Aspirin from Salicylic Acid & Acetic Acid is an equilibrium process. We want to make as much aspirin as possible.

Decrease in conc. of a component: Remove Aspirin


$K_{eq}$  must remain constant

$$K_{eq} = \frac{[\text{Asp.}]}{[\text{AA}][\text{SA}]}$$

To keep  $K_{eq}$  constant, we need to decrease the denominator – reaction will shift toward the products (use up AA & SA, & make more aspirin).

If we remove aspirin, the numerator gets smaller

# Changing Components, Concentration, or Partial Pressure



$$K_{eq} = \frac{[\text{Asp.}]}{[\text{AA}][\text{SA}]}$$

Synthesis of Aspirin from Salicylic Acid & Acetic Acid is an equilibrium process. We want to make as much aspirin as possible.

What if we added aspirin?

$K_{eq}$  must remain constant

$$K_{eq} = \frac{[\text{Asp.}]}{[\text{AA}][\text{SA}]}$$

To keep  $K_{eq}$  constant, we need to increase the denominator – reaction will shift toward the reactants (make more AA & SA).

If we add aspirin, the numerator gets larger

# Changing Components, Concentration, or Partial Pressure

For the following reaction,




what happens to the equilibrium if:

(a)  $\text{O}_2$  is added to the system?


(b)  $\text{NO}$  is removed?

# Changing Volume or External Pressure



**When pressure is increased:**


Equilibrium shifts to produce smaller # moles of gas



Toward products – 2 moles < 4 moles

**When Pressure is decreased:**

Equilibrium shifts to produce larger # moles of gas



Toward reactants – 4 moles > 2 moles

**If a reaction has the same number of moles of gas on both sides** (e.g.  $\text{N}_2(\text{g}) + \text{O}_2(\text{g}) \rightleftharpoons 2\text{NO}(\text{g})$ ):

Changes in external pressure do not affect equilibrium

# Changing Volume or External Pressure

For the following reaction,



what happens to the equilibrium if:

(a) The volume of the system is increased?

(b) The external pressure exerted on the system is increased?





# Adding a Catalyst or Inhibitor


## Catalysts:

- Catalysts lower activation energy.
- Cause a reaction to reach equilibrium faster
- DO NOT AFFECT THE EQUILIBRIUM ITSELF

## Inhibitors

- Act to slow the rate of a reaction
- Prevent equilibrium from being reached as quickly
- DO NOT AFFECT THE EQUILIBRIUM ITSELF
- It may seem like the reaction is not occurring at all, but in reality it is just infinitely slow.

## Which way does the reaction shift if.....



NO is added?


Volume of reaction vessel is reduced?

The total internal pressure is increased by adding He gas?

The temperature is increased?

A catalyst is added?

# How does each of the following changes affect the yield of NO at equilibrium?



Increase  $[\text{NH}_3]$

Increase  $[\text{H}_2\text{O}]$

Decrease  $[\text{O}_2]$

Decrease the volume of the container

Add a catalyst

Increase temperature