



# Chapter 17

## Acids & Bases



# Some Polyatomic Ions that are Important for Acids & Bases

Ammonium	$\text{NH}_4^+$	Nitrate	$\text{NO}_3^-$
Hydronium	$\text{H}_3\text{O}^+$	Nitrite	$\text{NO}_2^-$
Acetate	$\text{CH}_3\text{COO}^-$	Phosphate	$\text{PO}_4^{3-}$
Carbonate	$\text{CO}_3^{2-}$	Perchlorate	$\text{ClO}_4^-$
Hydroxide	$\text{OH}^-$	Sulfate	$\text{SO}_4^{2-}$
Cyanide	$\text{CN}^-$		


You should know these ions

# What are Acids & Bases?

## Arrhenius Definition


### Acid:

A substance that, when dissolved in water, increases the concentration of hydrogen ( $\text{H}^+$ ) ions (aka protons).



### Base:

A substance that, when dissolved in water, increases the concentration of hydroxide ions ( $\text{OH}^-$ ).




# What are Acids & Bases?

## Brønsted–Lowry Definition

### Acid:

A proton ( $\text{H}^+$ ) donor


- Must have a removable proton
- Proton goes to a base



### Base:

A proton ( $\text{H}^+$ ) acceptor

- Must have a pair of non-bonding electrons



# Lewis Acids & Bases

## Acid:



An electron pair acceptor

- Do not need to have a removable proton
- Must have an empty orbital to put electrons in

## Base:

An electron pair donor

- Must have a pair of non-bonding electrons



# Strength of Acids & Bases

## Strong Acids & Bases: Complete dissociation

- Non  $\text{H}^+$ / $\text{OH}^-$  ions form spectator ions
- Can use basic stoichiometry (CHM 101) in calculations
- No undissociated reactant left in solution

## Weak Acids & Bases: Incomplete dissociation

- Equilibrium process
- Equilibrium constants are  $K_a$  or  $K_b$
- Titrations involve stoichiometry and equilibrium calcs.

## Aqueous Solutions

- Acid or Base reacts with water
  - Water acts as a weak acid or base in the reaction

# Factors impacting acid strength

**Basic concept: easier to lose H, stronger acid**

Charge:

- Higher charge = stronger bond to  $\text{H}^+$  = weaker acid
- Ex:  $\text{HAsO}_4^{2-}$  weaker than  $\text{H}_2\text{AsO}_4^-$  weaker than  $\text{H}_3\text{AsO}_4$

Bond length:

- Shorter bonds are stronger so  $\text{H}^+$  harder to remove
- Ex: HF weaker than HCl weaker than HBr

Electronegativity:



- Less electronegative, share  $e^-$  more equally, stronger bond
- Ex: P less electronegative than S,  $\text{H}_3\text{PO}_4$  weaker than  $\text{H}_2\text{SO}_4$

For Oxoacids

- If all else is equal, fewer double bonds = weaker acid
- Ex:  $\text{H}_2\text{SO}_3$  weaker than  $\text{H}_2\text{SO}_4$
- More double bonds = more resonance = more stability with  $\text{H}^+$  removed

Stability of conjugate base: more stable anion = more acidic

# Binary Acid Trends in the Periodic Table

4A	5A	6A	7A	
<b>CH<sub>4</sub></b> Neither acid nor base	<b>NH<sub>3</sub></b> Weak base $K_b = 1.8 \times 10^{-5}$	<b>H<sub>2</sub>O</b>	<b>HF</b> Weak acid $K_a = 6.8 \times 10^{-4}$	 Increasing acid strength
<b>SiH<sub>4</sub></b> Neither acid nor base	<b>PH<sub>3</sub></b> Very weak base $K_b = 4 \times 10^{-28}$	<b>H<sub>2</sub>S</b> Weak acid $K_a = 9.5 \times 10^{-8}$	<b>HCl</b> Strong acid	
		<b>H<sub>2</sub>Se</b> Weak acid $K_a = 1.3 \times 10^{-4}$	<b>HBr</b> Strong acid	
 Increasing acid strength				

For binary acids:

Acidity increases left to right across a row

- Electronegativity increases left to right

Acidity increases top to bottom down a group

- Atomic size increases down a group



# Common Acids & Bases You Will Need to Know

## Strong Acids:

Hydrochloric Acid	HCl
Sulfuric Acid	H <sub>2</sub> SO <sub>4</sub>
Nitric Acid	HNO <sub>3</sub>
Perchloric Acid	HClO <sub>4</sub>
Hydrobromic Acid	HBr
Hydroiodic Acid	HI

## Strong Bases:

Soluble Hydroxides:

Sodium	NaOH
Potassium	KOH
Lithium	LiOH
Barium	Ba(OH) <sub>2</sub>
etc.	

## Weak Acids:

Carbonic Acid	H <sub>2</sub> CO <sub>3</sub>
Phosphoric Acid	H <sub>3</sub> PO <sub>4</sub>
Acetic Acid	CH <sub>3</sub> COOH
Hydrofluoric Acid	HF
Nitrous Acid	HNO <sub>2</sub>
Hydrocyanic acid	HCN
Carboxylic Acids	R-COOH

## Weak Bases:

Ammonia	NH <sub>3</sub>
Amines	R-NH <sub>2</sub> , R <sub>2</sub> NH
Insoluble/slightly soluble hydroxides	


# Organic Acids: Carboxylic Acids (–COOH)

## Weak organic acids


- –COOH group on molecule is acidic
- Removal of proton ( $\text{H}^+$ ) creates resonance structure
- Stabilizes anion

## Never fully dissociate in water


- Equilibrium process




## Examples:




Formic Acid



Acetic Acid



Butyric Acid



Benzoic Acid


# Organic Bases: Amines (contain N)

## Weak organic bases


- Derivatives of ammonia
- N has lone pair of electrons to accept a proton

Also do not fully dissociate in water


- Equilibrium process




## Examples:



Methyl amine



Dimethyl amine



Trimethyl amine

# Organic Acids & Bases: Things to keep in mind

## Organic Acids & Bases:

Only acidic proton

Acid:  $\text{CH}_3\text{COOH}$  Becomes  $\text{CH}_3\text{COO}^-$



Amine:  $\text{CH}_3\text{NH}_2$  Becomes  $\text{CH}_3\text{NH}_3^+$

Carbon: no lone pairs

- Cannot accept extra protons
- Protons bonded to carbon are not acidic

Added proton bonds via lone pair on N

# Conjugate Acid–Base Pairs




**Conjugate Acid:** Formed from the **base** after  $\text{H}^+$  is added

**Conjugate Base:** Formed from the **acid** after  $\text{H}^+$  is lost



Each acid has a conjugate base, each base has a conjugate acid. For substances that can both gain or lose a proton, whether they act as an acid or base depends on the system.

# Conjugate Acid–Base Pair Examples

1. Give the conjugate base of each of the following acids:




2. Write the formula for the conjugate acid of each of the following bases:





# H<sup>+</sup> Ion in Water

**H<sup>+</sup> is simply a proton – an H atom with no electron**


- In water, clusters of hydrated H<sup>+</sup> form
- Simplest cluster is the hydronium ion: H<sub>3</sub>O<sup>+</sup>



- H<sup>+</sup> (aq) & H<sub>3</sub>O<sup>+</sup>(aq) are used interchangeably




# Proton Transfer Reactions: Aqueous Acid



- **HCl** (the BL acid) donates a proton (H<sup>+</sup>)
- **Water** (the BL base) accepts the proton
- The conjugate base of the acid (Cl<sup>-</sup>) and the conjugate acid of the base (H<sub>3</sub>O<sup>+</sup>) are formed




# Proton Transfer Reactions: Aqueous Base



- **Water** (the BL acid) donates a proton (H<sup>+</sup>)
- **Ammonia** (the BL base) accepts the proton
- Water is **AMPHIPROTIC** – it can act as either an acid or a base (donate or accept a proton)

# Proton Transfer Reactions: Non-Aqueous




- **HCl** (the BL acid) donates a proton ( $\text{H}^+$ )
- **Ammonia** (the BL base) accepts the proton
- Can occur in the gas phase – water not needed
- Advantage of Brønsted–Lowry definition over Arrhenius definition
- Lewis definition even more broad (electron pair donor/acceptor)

# Acid–Base Properties of Water: Autoionization

**Autoionization:** In pure water, one water molecule can donate a proton to another water molecule


- Essentially the water ionizes itself (“auto”)




This is why pure water can conduct electricity

# Autoionization: An Equilibrium Process

Consider the autoionization of water at 25°C



$$[\text{H}_3\text{O}^+] = [\text{OH}^-] = 1.0 \times 10^{-7} \text{M}$$



This  $\text{H}_3\text{O}^+$  &  $\text{OH}^-$  concentration is where the pH of 7 for pure water comes from

$$K_w = [\text{H}_3\text{O}^+][\text{OH}^-] = 1.0 \times 10^{-14} \text{ (ion-product constant)}$$

$K_w$  is very small = favors reactants ( $\text{H}_2\text{O}$ )

$K_w$  applies to both pure water and aqueous solutions

- If know acid concentration, can use  $K_w$  to find the base concentration & vice versa

# pH & pOH

## Method of Measuring Acidity

- Power of the Hydrogen Ion

### Formulas:

- $\text{pH} = -\log[\text{H}_3\text{O}^+]$
- $[\text{H}_3\text{O}^+] = 10^{(-\text{pH})}$
- $\text{pOH} = -\log[\text{OH}^-]$
- $[\text{OH}^-] = 10^{(-\text{pOH})}$
- $K_w = [\text{H}_3\text{O}^+][\text{OH}^-] = 1.0 \times 10^{-14} \text{M}$
- $\text{p}K_w = \text{pH} + \text{pOH} = 14$

Neutral:  $[\text{H}_3\text{O}^+] = [\text{OH}^-]$   $\text{pH} = 7$

Acidic:  $[\text{H}_3\text{O}^+] > [\text{OH}^-]$   $\text{pH} < 7$

Basic:  $[\text{H}_3\text{O}^+] < [\text{OH}^-]$   $\text{pH} > 7$

Sample	pH Value
Gastric juice in the stomach	1.0–2.0
Lemon juice	2.4
Vinegar	3.0
Grapefruit juice	3.2
Orange juice	3.5
Urine	4.8–7.5
Water exposed to air*	5.5
Saliva	6.4–6.9
Milk	6.5
Pure water	7.0
Blood	7.35–7.45
Tears	7.4
Milk of magnesia	10.6
Household ammonia	11.5

Sig Figs: # sig figs in concentration = # sig figs after decimal point in pH/pOH

# What is log?

## Consider the number $1.0 \times 10^{-3}$

- Log refers to base 10
- Essentially, it refers to the exponent in a number written in scientific notation
- It tells you the magnitude (size) of the number
- The log of  $1.0 \times 10^{-3}$  is  $-3$
- The formula for pH is  $-\log$  to eliminate the negative sign in the answer

## Consider the number $2.8 \times 10^{-3}$

- Log still refers primarily to the exponent, but the actual value is impacted by the rest of the number
- The log of  $2.8 \times 10^{-3}$  will be close to, but not exactly,  $-3$
- $\text{Log}(2.8 \times 10^{-3}) = -2.6$

Low pH values are acidic because concentrations generally have negative exponents.  $1 \times 10^{-3} \text{M} > 1 \times 10^{-10} \text{M}$

# Measuring pH

## Most Accurate: pH meter

- Measures the voltage in a solution to determine concentration & pH



## Other methods:

- Litmus paper
  - Red litmus paper turns blue above ~ pH 8
  - Blue litmus paper turns red below ~ pH 5
- Indicators
  - In solution or on pHYdrion paper

	pH range for color change													
	0	2	4	6	8	10	12	14						
Methyl violet	Yellow													Violet
Thymol blue	Red													Blue
Methyl orange														
Methyl red														
Bromthymol blue														
Phenolphthalein														
Alizarin yellow R														

# Concentrated vs. Dilute Solutions

Do we need to take the autoionization of water into account when determining pH?

## Example 1: Concentrated Solutions

Consider an aqueous 0.010M solution of nitric acid.

Two reactions are occurring:



The  $[\text{H}_3\text{O}^+]$  from ionization of water is negligible:

$$0.010\text{M} + 0.0000001\text{M} = 0.0100001\text{M}$$

It can be ignored




# Concentrated vs. Dilute Solutions

## Example 2: Dilute Solutions

Consider an aqueous  $1.0 \times 10^{-6} \text{M}$  solution of nitric acid.

Two reactions are again occurring:



\*Likely somewhat less due to Le Châtelier's Principle

The  $[\text{H}_3\text{O}^+]$  from ionization of water is 10% of the amount contributed by the acid:

$$1.0 \times 10^{-6} \text{M} + 0.1 \times 10^{-6} \text{M} = 1.1 \times 10^{-6} \text{M}$$

It CANNOT be ignored

Contribution from autoionization of water must be taken into account if acid/base provides  $< 10^{-6} \text{M}$   $\text{H}_3\text{O}^+/\text{OH}^-$

# pH Calculations for Strong Acids/Bases

1. Calculate  $[H^+]$  at  $25^\circ C$  for an aqueous solution in which  $[OH^-] = 0.00045M$ . Indicate whether it is acidic, basic, or neutral. A:  $2.2 \times 10^{-11}M$ ; basic
2. Find the pH and pOH of a  $0.0050M$  HBr solution at  $25^\circ C$   
pH: 2.30; pOH: 11.70
3. Calculate the  $H_3O^+$  and  $OH^-$  concentrations at  $25^\circ C$  of an aqueous  $0.010M$  solution of nitric acid.  $[H_3O^+]$ :  $0.010M$   
 $[OH^-]$ :  $1.0 \times 10^{-12}$



5. Calculate the pH made from 15.00mL of 1.00M HCl diluted to 0.500L. A:1.523

6. What is the concentration of a solution of  $\text{Ba}(\text{OH})_2$  for which the pH is 10.05? A:  $5.6 \times 10^{-5}\text{M}$

# Strength of Acids & Bases

## Strong Acids & Bases: Complete dissociation



- Strong electrolytes
- Good conductors of electricity
- Completely ionized in aqueous solution; no original compound remains
- Conjugate has no measurable strength
- Single arrow – not equilibrium





# Strength of Acids & Bases

## Weak Acids & Bases: Incomplete dissociation



- Some of original compound remains along with ions
- Equilibrium process; represented by double arrow
- Dissociation is governed by an equilibrium constant
  - $K_a$  or  $K_b$
- Poor conductors of electricity
- Conjugates can act as acids/bases



# Strong Acid (HCl)




# Weak Acid (HF)



Strong vs. weak  
simulation

## 2 Factors affect pH of solution:

- nature of acid/base
- concentration




# Relative Strengths of Conjugate Acid–Base Pairs

Strong Acids/Bases give weak conjugates and vice versa

Table 16.2 Relative Strengths of Conjugate Acid-Base Pairs		
	Acid	Conjugate Base
<div style="display: flex; flex-direction: column; align-items: center;"> <div style="margin-bottom: 20px;">↑ Acid strength increases</div> <div style="margin-bottom: 20px;">Strong acids</div> <div style="margin-bottom: 20px;">Weak acids</div> <div>↓</div> </div>	HClO <sub>4</sub> (perchloric acid)	ClO <sub>4</sub> <sup>-</sup> (perchlorate ion)
	HI (hydroiodic acid)	I <sup>-</sup> (iodide ion)
	HBr (hydrobromic acid)	Br <sup>-</sup> (bromide ion)
	HCl (hydrochloric acid)	Cl <sup>-</sup> (chloride ion)
	H <sub>2</sub> SO <sub>4</sub> (sulfuric acid)	HSO <sub>4</sub> <sup>-</sup> (hydrogen sulfate ion)
	HNO <sub>3</sub> (nitric acid)	NO <sub>3</sub> <sup>-</sup> (nitrate ion)
	H <sub>3</sub> O <sup>+</sup> (hydronium ion)	H <sub>2</sub> O (water)
	HSO <sub>4</sub> <sup>-</sup> (hydrogen sulfate ion)	SO <sub>4</sub> <sup>2-</sup> (sulfate ion)
	HF (hydrofluoric acid)	F <sup>-</sup> (fluoride ion)
	HNO <sub>2</sub> (nitrous acid)	NO <sub>2</sub> <sup>-</sup> (nitrite ion)
	HCOOH (formic acid)	HCOO <sup>-</sup> (formate ion)
	CH <sub>3</sub> COOH (acetic acid)	CH <sub>3</sub> COO <sup>-</sup> (acetate ion)
	NH <sub>4</sub> <sup>+</sup> (ammonium ion)	NH <sub>3</sub> (ammonia)
	HCN (hydrocyanic acid)	CN <sup>-</sup> (cyanide ion)
	H <sub>2</sub> O (water)	OH <sup>-</sup> (hydroxide ion)
	NH <sub>3</sub> (ammonia)	NH <sub>2</sub> <sup>-</sup> (amide ion)
		<div style="display: flex; flex-direction: column; align-items: center;"> <div>↓ Base strength increases</div> </div>

Stronger acids will dominate over weaker acids






# Acid Ionization Constant: $K_a$

## Equilibrium constant for acid dissociation

Dissociation of acid in water:



$$K_{\text{eq}} = \frac{[\text{H}_3\text{O}^+][\text{A}^-]}{[\text{HA}]} = K_a$$

$K_a$  is the acid ionization (aka acid dissociation) constant






- Quantitative measure of acid strength
- Higher  $K_a$  = stronger acid
- Sometimes discussed in terms of  $\text{p}K_a$

$$\text{p}K_a = -\log K_a$$

$K_a$  ↑ Weak acid strength ↑

$\text{p}K_a$  ↑ Weak acid strength ↓

# K<sub>a</sub> Values for Some Common Weak Acids

Name of Acid	Formula	Structure	K <sub>a</sub>	Conjugate Base	K <sub>b</sub>
Hydrofluoric acid	HF	H—F	$7.1 \times 10^{-4}$	F <sup>-</sup>	$1.4 \times 10^{-11}$
Nitrous acid	HNO <sub>2</sub>	O=N—O—H	$4.5 \times 10^{-4}$	NO <sub>2</sub> <sup>-</sup>	$2.2 \times 10^{-11}$
Acetylsalicylic acid (aspirin)	C <sub>9</sub> H <sub>8</sub> O <sub>4</sub>		$3.0 \times 10^{-4}$	C <sub>9</sub> H <sub>7</sub> O <sub>4</sub> <sup>-</sup>	$3.3 \times 10^{-11}$
Formic acid	HCOOH	H—C(=O)—O—H	$1.7 \times 10^{-4}$	HCOO <sup>-</sup>	$5.9 \times 10^{-11}$
Ascorbic acid*	C <sub>6</sub> H <sub>8</sub> O <sub>6</sub>		$8.0 \times 10^{-5}$	C <sub>6</sub> H <sub>7</sub> O <sub>6</sub> <sup>-</sup>	$1.3 \times 10^{-10}$
Benzoic acid	C <sub>6</sub> H <sub>5</sub> COOH		$6.5 \times 10^{-5}$	C <sub>6</sub> H <sub>5</sub> COO <sup>-</sup>	$1.5 \times 10^{-10}$
Acetic acid	CH <sub>3</sub> COOH		$1.8 \times 10^{-5}$	CH <sub>3</sub> COO <sup>-</sup>	$5.6 \times 10^{-10}$
Hydrocyanic acid	HCN	H—C≡N	$4.9 \times 10^{-10}$	CN <sup>-</sup>	$2.0 \times 10^{-5}$
Phenol	C <sub>6</sub> H <sub>5</sub> OH		$1.3 \times 10^{-10}$	C <sub>6</sub> H <sub>5</sub> O <sup>-</sup>	$7.7 \times 10^{-5}$

## Calculations Using $K_a$

1.) 0.100mol of HF is dissolved in 1.00L of water at 25°C. The pH at equilibrium was found to be 2.08. Calculate  $K_a$  and pKa.

**Note: Do not use approximation method if you have the value of X!**

A:  $K_a = 7.5 \times 10^{-4}$   
 $pK_a = 3.12$

2.) What is the pH of a weak 0.122M monoprotic acid whose  $K_a$  is  $5.7 \times 10^{-6}$ ?

A: pH = 3.08

3.) Find the pH of a 0.20M solution of Formic Acid ( $\text{HCHO}_2$ )

Look up  $K_a$  in table:

# Determining Relative Acidity

1.) Which is the stronger acid, HF or HNO<sub>2</sub>?

Look up  $K_a$  in table:


HF:  $7.5 \times 10^{-4}$

HNO<sub>2</sub>:  $4.6 \times 10^{-4}$

# Percent Ionization and $K_a$

- Indicates the extent to which an acid is ionized
- Only matters for weak acids (100% for strong)
- Higher percent ionization = more ions = stronger acid
  - Ex: 0.1M HCl, pH = 1.00
  - 0.1M HCOOH, pH = 2.38


$$\text{Percent Ionization} = \frac{[\text{H}_3\text{O}^+]_{\text{eq}}}{[\text{HA}]_{\text{initial}}} \times 100$$



% ionization of a weak acid decreases as the initial concentration of the acid increases

# Calculations Using Percent Ionization

1.) Niacin, one of the B vitamins, has the following structure:



a.) If a 0.020M solution has a pH of 3.26, what is the  $K_a$  for niacin?  
A:  $1.6 \times 10^{-5}$

b.) What is the percent ionization of the 0.020M solution?




2.) A 0.0100M solution of  $\text{HNO}_2$  is 19% ionized at equilibrium. What is the  $K_a$ ?

# Polyprotic Acids

## Acids that have more than one ionizable proton

- Ionize in successive steps



- Each step has its own  $K_a$
- Easier to remove the first proton than the second, etc.
  - $K_{a1} > K_{a2} > K_{a3} \dots$
- Successive equilibrium constants have less and less impact on pH
  - May not need to use all  $K$  values to solve a problem

# Polyprotic Acids

If the difference between the  $K_a$  values for the first and 2<sup>nd</sup>, 3<sup>rd</sup>, etc.  $K_a$  values is  $10^3$  or more, the pH generally depends on only the first dissociation.

Name	Formula	$K_{a1}$	$K_{a2}$	$K_{a3}$
Ascorbic	$\text{H}_2\text{C}_6\text{H}_6\text{O}_6$	$8.0 \times 10^{-3}$	$1.6 \times 10^{-12}$	
Carbonic	$\text{H}_2\text{CO}_3$	$4.3 \times 10^{-7}$	$5.6 \times 10^{-11}$	
Citric	$\text{H}_3\text{C}_6\text{H}_5\text{O}_7$	$7.4 \times 10^{-4}$	$1.7 \times 10^{-5}$	$4.0 \times 10^{-7}$
Oxalic	$\text{H}_2\text{C}_2\text{O}_4$	$5.9 \times 10^{-2}$	$6.4 \times 10^{-5}$	
Phosphoric	$\text{H}_3\text{PO}_4$	$7.5 \times 10^{-3}$	$6.2 \times 10^{-8}$	$4.2 \times 10^{-13}$
Sulfurous	$\text{H}_2\text{SO}_3$	$1.7 \times 10^{-2}$	$6.4 \times 10^{-8}$	
Sulfuric	$\text{H}_2\text{SO}_4$	Large	$1.2 \times 10^{-2}$	
Tartaric	$\text{H}_2\text{C}_4\text{H}_4\text{O}_6$	$1.0 \times 10^{-3}$	$4.6 \times 10^{-5}$	

# Polyprotic Acid Calculations

Calculate the pH of a 1.5M solution of tartaric acid.


$$K_{a1} = 1.0 \times 10^{-3} \quad K_{a2} = 4.6 \times 10^{-5}$$

A: pH=

# Base Ionization Constant: $K_b$

## Equilibrium constant for base dissociation

Weak bases react with water to produce hydroxide ions:



$$K_{eq} = \frac{[HB^+][OH^-]}{[B]} = K_b$$

$K_b$  is the base ionization constant

- Quantitative measure of base strength
- Higher  $K_b$  = stronger base
- Sometimes discussed in terms of  $pK_b$

$$pK_b = -\log K_b$$








$K_b$  ↑ Weak base strength ↑

$pK_b$  ↑ Weak base strength ↓

# Types of Weak Bases


- Neutral substances that have an atom with a non-bonding pair of electrons
  - Examples include ammonia ( $\text{NH}_3$ ) & amines
  - Amines described earlier in the chapter (slide 11)
- Anions (conjugate bases) of weak acids
  - Examples include  $\text{HCO}_3^-$ ,  $\text{ClO}^-$ ,  $\text{HS}^-$
  - Act as  $\text{H}^+$  acceptors in water
  - Anions of strong acids do not act as bases – not an equilibrium process
- Insoluble/slightly soluble hydroxides
  - Soluble hydroxides are strong bases
  - $\text{Ca}(\text{OH})_2$ ,  $\text{Fe}(\text{OH})_3$
  - Solubility equilibria ( $K_{\text{sp}}$ ) will be discussed in the next chapter

# K<sub>b</sub> Values for Some Common Weak Bases


Name of Base	Formula	Structure	K <sub>b</sub> *	Conjugate Acid	K <sub>a</sub>
Ethylamine	C <sub>2</sub> H <sub>5</sub> NH <sub>2</sub>		$5.6 \times 10^{-4}$	C <sub>2</sub> H <sub>5</sub> NH <sub>3</sub> <sup>+</sup>	$1.8 \times 10^{-11}$
Methylamine	CH <sub>3</sub> NH <sub>2</sub>		$4.4 \times 10^{-4}$	CH <sub>3</sub> NH <sub>3</sub> <sup>+</sup>	$2.3 \times 10^{-11}$
Ammonia	NH <sub>3</sub>		$1.8 \times 10^{-5}$	NH <sub>4</sub> <sup>+</sup>	$5.6 \times 10^{-10}$
Pyridine	C <sub>5</sub> H <sub>5</sub> N		$1.7 \times 10^{-9}$	C <sub>5</sub> H <sub>5</sub> NH <sup>+</sup>	$5.9 \times 10^{-6}$
Aniline	C <sub>6</sub> H <sub>5</sub> NH <sub>2</sub>		$3.8 \times 10^{-10}$	C <sub>6</sub> H <sub>5</sub> NH <sub>3</sub> <sup>+</sup>	$2.6 \times 10^{-5}$
Caffeine	C <sub>8</sub> H <sub>10</sub> N <sub>4</sub> O <sub>2</sub>		$5.3 \times 10^{-14}$	C <sub>8</sub> H <sub>11</sub> N <sub>4</sub> O <sub>2</sub> <sup>+</sup>	0.19
Urea	(NH <sub>2</sub> ) <sub>2</sub> CO		$1.5 \times 10^{-14}$	H <sub>2</sub> NCONH <sub>3</sub> <sup>+</sup>	0.67

# $K_a$ & $K_b$ Relationship for Conjugate Acid/Base Pairs in Water


## Base & water



## Conjugate acid & water



Add the two reactions together: multiply K values (Ch15)



$$K_w = K_a \times K_b = 1.0 \times 10^{-14}$$

$$1.8 \times 10^{-5} \times 5.6 \times 10^{-10} = 1.0 \times 10^{-14}$$

$$\text{p}K_a + \text{p}K_b = 14$$
$$9.26 + 4.74 = 14$$

High  $K_a$  must have  
Low  $K_b$  & vice versa



## Calculations Using $K_b$

1.) What is the pH of a 0.15M solution of  $\text{NH}_3$ ?  $K_b = 1.8 \times 10^{-5}$

A: pH = 11.22 49

2.) Codeine is a weak organic base. A  $5.0 \times 10^{-3} \text{M}$  solution of codeine has a pH of 9.95. Calculate the  $\text{pK}_b$  and the value of  $K_b$  for this base.

**Reminder: Do not use approximation method if you have the value of X!**

A:  $K_b = 1.62 \times 10^{-6}$   
 $\text{pK}_b = 5.79$

3.) A solution of ammonia in water has a pH of 11.17.  
What is the molarity of the solution?  $K_b = 1.8 \times 10^{-5}$


# Acid–Base Properties of Salts

Salts are ionic compounds – contain cation (+) & anion (–)

- Can be formed from acid/base neutralization
- Dissociate (separate into ions) when dissolve in water
- Can sometimes impact pH through hydrolysis


## Hydrolysis:

- Hydro = water
- lysis = to cut
- Reactions that break water into  $\text{H}^+$  ( $\text{H}_3\text{O}^+$ ) &  $\text{OH}^-$



**Extent to which a salt causes hydrolysis determines extent of its acid–base properties (if any)**

## Reminders: Strong vs. Weak No Equilibrium vs. Equilibrium





# Acid-Base Properties of Salts

Hydrolysis can be caused by both cations & anions,  
but does not happen in every situation

Salts formed from conjugates of a strong acid + a strong base do not cause hydrolysis

For salt solutions: no hydrolysis = neutral (no  $\text{OH}^-$  or  $\text{H}_3\text{O}^+$ )



# Acid–Base Properties of Salts


Soluble ions derived from a weak base or a weak acid will cause hydrolysis

Ex 1: **NaOH: strong base**

**CH<sub>3</sub>COOH: weak acid**



Weak base




Ex 2: **NH<sub>3</sub>: weak base**

**HCl: strong acid**



Weak acid




# What if an ion can act as either an acid or a base?

Occurs with salts from polyprotic acids such as  $\text{H}_2\text{CO}_3$ ,  $\text{H}_3\text{PO}_4$ ,  $\text{H}_2\text{SO}_4$ .


Need to compare  $K_a$  and  $K_b$  values

- Example:  $\text{NaHCO}_3$

Acting  
as acid



Acting  
as base



- $K_b > K_a$  so solution will be basic!




# Determining if a Salt Solution is Acidic, Basic, or Neutral

Look at the ions that the salt dissociates into:

- If the base that would contain the cation is strong – not acidic
- If the acid that would contain the anion is strong – not basic
- If both cation & anion have a strong parent – neutral
- If the base that would contain the cation is weak – acidic
- If the acid that would contain the anion is strong – basic
- If both cation & anion have a weak parent
  - Acidity/basicity depends on relative strength
  - Math can be complicated
  - Qualitatively:
    - $K_a < K_b$  basic solution
    - $K_a > K_b$  acidic solution
    - $K_a \approx K_b$  pretty close to neutral

# Determining if a Salt Solution is Acidic, Basic, or Neutral

1.) Predict whether aqueous solutions of the following compounds are acidic, basic, or neutral.



2.) For each of the following pairs, indicate which salt would form the more acidic (i.e. less basic) 0.010M solution.

a.)  $\text{NaNO}_3$  or  $\text{Fe}(\text{NO}_3)_3$

b.)  $\text{CH}_3\text{NH}_3\text{Cl}$  or  $\text{BaCl}_2$

c.)  $\text{KNO}_2$  or  $\text{KNO}_3$

d.)  $(\text{NH}_4)_2\text{SO}_4$  or  $(\text{NH}_4)_2\text{SO}_3$

3.) Calculate the  $[\text{OH}^-]$  and pH of a 0.10M NaCN solution.  $K_a$  for HCN is  $4.9 \times 10^{-10}$ .

A:  $[\text{OH}^-] = 1.4 \times 10^{-3}\text{M}$   
pH = 11.16

4.) Calculate the pH of a 0.42M  $\text{NH}_4\text{Cl}$  solution.  
 $K_b$  for  $\text{NH}_3$  is  $1.8 \times 10^{-5}$


A: pH = 4.82

# Hydration of Metal Ions

Salts with small, highly charged metal cations (e.g.  $\text{Al}^{3+}$ ,  $\text{Fe}^{3+}$ ,  $\text{Cr}^{3+}$ ,  $\text{Be}^{2+}$ ) and the conjugate base of a strong acid can also have acidic properties

Example:

$\text{AlCl}_3$  dissolved in water produces:



- $\text{e}^-$  in water molecules are pulled toward metal ion
- O–H bonds in attached waters become more polarized
- $\text{H}^+$  dissociates from water molecule forming  $\text{H}_3\text{O}^+$

# Hydration of Metal Ions



Higher charge & smaller size make cations more acidic



# Hydration of Metal Ions

Which member of each pair would produce a more acidic solution & why?

1.)  $\text{CuCl}$  or  $\text{Cu}(\text{NO}_3)_2$

2.)  $\text{CrCl}_3$  or  $\text{NiCl}_3$